

REVIEW ARTICLE

# Vitamin B<sub>12</sub> – a scoping review for Nordic Nutrition Recommendations 2023

Anne-Lise Bjørke-Monsen<sup>1\*</sup> and Vegard Lysne<sup>2</sup>

<sup>1</sup>Department of Medical Biochemistry and Pharmacology, Haukeland University Hospital, Bergen, Norway;

<sup>2</sup>Norwegian Institute of Public Health, Oslo, Norway

## Popular scientific summary

- Vitamin B<sub>12</sub> (cobalamin) refers to a group of cobalt-containing compounds, corrinoids, that are biologically active in humans.
- Vitamin B<sub>12</sub> is essential for normal development, metabolic and neurological function, and blood formation.
- Vitamin B<sub>12</sub> only occurs in animal foods or fortified foods.
- Since biomarkers for vitamin B<sub>12</sub> status have different limitations, recommendations should be based on a combination of biomarkers.
- There is no consensus regarding what levels of vitamin B<sub>12</sub> biomarkers should be considered optimal.

## Abstract

Vitamin B<sub>12</sub> (cobalamin) is essential for normal metabolic function, and even moderate deficiency of this vitamin has negative health effects. Vitamin B<sub>12</sub> is found in animal foods, and as vegetarian diets are increasingly popular in Western countries, one might expect a higher prevalence of vitamin B<sub>12</sub> deficiency in the Nordic population. Setting recommendations for vitamin B<sub>12</sub> intake has proven to be difficult, as uptake of vitamin B<sub>12</sub> varies substantially, the clinical deficiency symptoms are often diffuse, and there is no clear agreement on the decision limits for vitamin B<sub>12</sub> deficiency. Vitamin B<sub>12</sub> deficiency is reported to be particularly common among pregnant women and infants, despite the fact that less than 1% of Norwegian pregnant women have a cobalamin intake below the Nordic Nutrition Recommendations 2012-recommended level of 2.0 µg/day. In addition, the assumption that breast milk contains sufficient vitamin B<sub>12</sub> for optimal health and neurodevelopment during the first 6 months of life does not comply with the high prevalence of insufficient vitamin B<sub>12</sub> status in this age group. Recommended intakes of vitamin B<sub>12</sub> vary among age groups and must be based on markers of cobalamin status, indicating an optimal intracellular biochemical status, and not merely absence of clinical signs of vitamin B<sub>12</sub> deficiency.

Keywords: *Vitamin B<sub>12</sub>, cobalamin; one-carbon metabolism; homocysteine; methylmalonic acid, nutrition recommendations*

Received: 12 February 2022; Revised: 2 August 2022; Accepted: 27 September 2023; Published: 08 November 2023

The aim of this scoping review is to describe the totality of evidence for the role of vitamin B<sub>12</sub> for health-related outcomes as a basis for setting and updating dietary reference values (DRVs) for the Nordic Nutrition Recommendations (NNR) 2023 (Box 1). Vitamin B<sub>12</sub> is the common term for a group of cobalt-containing compounds (corrinoids) that are biologically active in humans. Cobalamin is used synonymously with vitamin B<sub>12</sub>.

Vitamin B<sub>12</sub> has two metabolic functions. Methylcobalamin serves as cofactor for methionine synthase, involved in the

transfer of a methyl group from folate to homocysteine, regenerating methionine. Adenosylcobalamin is a cofactor for methylmalonyl coenzyme A mutase, involved in the metabolism of odd-chain fatty acids, amino acids, and cholesterol. Insufficient activity of these enzymes, due to vitamin B<sub>12</sub> deficiency, results in the accumulation of homocysteine and/or methylmalonic acid (MMA), which therefore are considered functional markers of vitamin B<sub>12</sub> status (1).

Vitamin B<sub>12</sub> is produced by soil bacteria and is found in animal food and soil-contaminated food (2).

Vegetarian and vegan diets are increasingly popular in Western countries, particularly among young women (3). Although these diets are considered nutritionally adequate and healthy when appropriately planned, specific nutritional knowledge is needed to achieve an optimal vegetarian diet with regard to vitamin B<sub>12</sub> intake and status. However, this is not always the case, even among well-educated people, and a higher prevalence of vitamin B<sub>12</sub> deficiency in the Nordic population might be expected (4).

Vitamin B<sub>12</sub> is essential for normal metabolic function, and even moderate deficiency of this vitamin has negative health effects, particularly in young infants, where vitamin B<sub>12</sub> insufficiency is associated with impaired neurodevelopment (5–7). Vitamin B<sub>12</sub> deficiency is reported to be common among Norwegian pregnant women (8) and infants (9). However, only 0.8% of Norwegian mothers in pregnancy week 18 had a vitamin B<sub>12</sub> intake below the recommended intake (RI) in the NNRs 2012 of 2.0 µg/day (10), indicating that the recommendations may be too low.

Setting recommendations for vitamin B<sub>12</sub> intake is difficult, and there are a number of uncertainties to consider. The absorption rate of vitamin B<sub>12</sub> varies from 30 to 90%, reported estimated body stores vary from 2 to 3 mg, and daily losses range from 1.4 to 5.1 µg. The vitamin B<sub>12</sub> intake needed just to compensate for daily losses is estimated to be in the range from 3.8 to 20.7 µg per day (11). In addition, there is no clear agreement on what levels of serum vitamin B<sub>12</sub>, plasma holotranscobalamin (holoTC), the metabolic markers plasma total homocysteine (tHcy), and serum MMA constitute an optimal vitamin B<sub>12</sub> status. It is however important to remember that vitamin B<sub>12</sub> is essential for normal metabolic function; as long as vitamin B<sub>12</sub> is taken orally, either in food or in supplements containing a moderate vitamin B<sub>12</sub> content, the body will actively regulate the uptake in ileum, and there is no evidence that intakes up to 100 µg/d from foods and supplements represent a health risk (12).

## Methods

This scoping review follows the protocol developed within the NNR2023 project (13). The sources of evidence used in the scoping review follow the eligibility criteria described previously (2).

The main literature search for this review was performed in MEDLINE on July 1st, 2022, with a following search string: (b12, vitamin[MeSH Terms] OR cobalamin[MeSH Terms]) AND review[Publication Type] AND ('2011'[Date – Publication] : '3000'[Date – Publication]) AND Humans[Filter]. The number of hits was 453. Based on the title and abstract, a total of 32 articles were picked up, of which 18 were considered as relevant (11, 14–30). Three additional articles were identified through a supplementary search (31–33). We also identified relevant literature for this scoping review via 'snowballing'/citation chasing that was relevant for the background information. No published articles were considered by the committee as a qualified systematic review (34), and a de novo NNR2023 systematic review was published in 2023 (35).

## Physiology

Vitamin B<sub>12</sub> refers to a group of cobalt-containing compounds (corrinoids) that are biologically active in humans. Corrinoids also include vitamin B<sub>12</sub> analogs, which are not biologically active, such as cobamides and cobinamides. The vitamin B<sub>12</sub> molecule consists of a cobalt-containing corrin ring, a 5,6-dimethylbenzimidazole base, a sugar, and an aminopropanol group. A functional ligand is covalently bound to the cobalt at the upper axial position, which can either be a methyl-, adenosyl-, cyano-, or hydroxyl group.

### *Absorption and intestinal transport*

Dietary vitamin B<sub>12</sub> is absorbed via an active receptor-mediated physiological process and via a less efficient passive diffusion pathway. In the acidic environment of the stomach, dietary protein-bound vitamin B<sub>12</sub> is cleaved from the protein by gastric enzymes and bound to salivary-derived haptocorrin (HC). Stimulated by food intake, intrinsic

### **Box 1.** The Nordic Nutrition Recommendations (NNR) 2023 project

- This paper is one of many scoping reviews commissioned as part of the Nordic Nutrition Recommendations 2023 (NNR2023) project (13).
- The papers are included in the extended NNR2023 report, but, for transparency, these scoping reviews are also published in Food & Nutrition Research.
- The scoping reviews have been peer reviewed by independent experts in the research field according to the standard procedures of the journal.
- The scoping reviews have also been subjected to public consultations (see report to be published by the NNR2023 project).
- The NNR2023 committee has served as the editorial board.
- While these papers are a main fundament, the NNR2023 committee has the sole responsibility for setting dietary reference values in the NNR2023 project.

factor (IF) is produced by the parietal cells in the stomach, and both IF and the haptocorrin-vitamin B<sub>12</sub> complex are transported to the small intestine. In the less acidic environment of the upper small intestine, vitamin B12 is released from HC, and the free vitamin B<sub>12</sub> binds to IF. While HC binds all corronoids, IF for all practical purposes only recognizes and binds biologically active cobalamins.

The IF-vitamin B<sub>12</sub> complex is transported to the distal ileum, where it binds to the cubam receptor and enters the enterocytes through endocytosis. The availability of cubam receptors is the rate-limiting factor for receptor-mediated absorption of cobalamin. It has been reported that the fractional absorption decreases with increasing dose (14). Bioavailability studies have demonstrated that while about 50% of a 1 µg oral dose is absorbed, only 5% of a 25 µg dose is retained. Furthermore, during the subsequent 4–6 h, active uptake is restricted due to regeneration of cubam receptors (14). In total, a maximum of ~2 µg vitamin B<sub>12</sub> can be absorbed from a meal through the active route. Additionally, approximately 1.2% of vitamin B<sub>12</sub> is absorbed passively (36). Normally, this accounts for a small proportion of total vitamin B<sub>12</sub> absorption, but becomes relevant if vitamin B12 is consumed in high doses (>50 µg). Bioavailability of vitamin B<sub>12</sub> from various foods, as assessed by whole-body retention or fecal excretion, ranges from about 20% up to 90% at single doses of 0.25 µg to 5 µg (37). It is estimated that approximately 50% of dietary vitamin B<sub>12</sub> is absorbed by healthy adults with normal gastric function (38, 39).

#### *Transport in blood and enterohepatic circulation*

Upon entering the enterocyte, IF is degraded, and free vitamin B<sub>12</sub> is released into the cytosol and transported into the circulation through multidrug resistance protein 1 (MRP1). In the circulation, vitamin B<sub>12</sub> is transported bound to either HC or transcobalamin (TC). About 70–90% of circulating vitamin B<sub>12</sub> is bound to HC, which can be taken up by the liver through specific receptors. About 20–30% of circulating vitamin B<sub>12</sub> is bound to TC (holoTC), which is responsible for delivering vitamin B<sub>12</sub> to all tissues through endocytosis facilitated by the transcobalamin receptor (TCR) (14). HC-bound vitamin B<sub>12</sub> has a half-life of several days compared to the half-life of about an hour for holoTC (40). HC-bound vitamin B<sub>12</sub> and vitamin B<sub>12</sub> analogues are excreted from the liver through bile and cleaved from HC in the intestines. While the analogues are excreted in feces, vitamin B<sub>12</sub> may be reabsorbed, referred to as the enterohepatic circulation.

#### *Intracellular metabolism*

Once bound to the TCR on the target cells membrane, the holoTC-TCR complex is taken up into the cell by endocytosis. Within the lysosome, vitamin B<sub>12</sub> is released, the TC

is degraded, and the TCR is recycled back to the cell surface. A series of vitamin B<sub>12</sub> processing proteins facilitates the transport out of the lysosome (CblF/CblJ), removal of the upper axial ligand (CblC/CblX), partitioning to the cytosolic or mitochondrial compartment (CblD), and the formation of the two active cobalamin cofactors methylcobalamin (CblE/CblG) and adenosylcobalamin (CblA/CblB/CblH/mut) (1).

#### *Molecular functions*

Vitamin B<sub>12</sub> is a cofactor for only two enzymes in the human metabolism (2). Methylcobalamin is a cofactor for methionine synthase – the enzyme that catalyzes the conversion of homocysteine to methionine. This reaction requires the transfer of a methylgroup from 5'-methyl tetrahydrofolate. Intracellular deficiency of folate and/or vitamin B<sub>12</sub> therefore results in increased plasma concentrations of tHcy. Adenosylcobalamin is a cofactor for methylmalonyl-CoA mutase in the isomerization of methylmalonyl-CoA to succinyl-CoA. Vitamin B<sub>12</sub> insufficiency therefore results in increased serum MMA concentrations (2).

#### *Specific groups*

##### *Infants, children, and adolescents*

The vitamin B<sub>12</sub> status in the new-born is a result of maternal status during pregnancy, gestational age, and birth weight. Infants who are born premature or with a low birth weight (LBW) or with a vitamin B<sub>12</sub> deficient mother are prone to develop vitamin B<sub>12</sub> deficiency during the first weeks of life, something that may impair neurodevelopment (5, 9). During the first weeks of life, there is a considerable decrease in serum cobalamin, accompanied by a marked increase in plasma tHcy and MMA (41–43). The lowest vitamin B<sub>12</sub> levels and the highest plasma tHcy and MMA levels in childhood are seen in infants 6 weeks to 6 months of age. In older children (>6 months), serum vitamin B<sub>12</sub> increases and peaks at 3–7 years and then decreases, and median plasma tHcy remains low (<6 µmol/L) and increases from the age of 7 (43, 44).

##### *Pregnancy and lactation*

All markers of vitamin B<sub>12</sub> status are reduced in pregnant compared to non-pregnant women (8). Serum vitamin B<sub>12</sub> decreases during pregnancy. This is accompanied by a gradual increase in the metabolic markers, indicative of a functional intracellular vitamin B<sub>12</sub> depletion, despite the fact that both plasma tHcy and MMA are substantially reduced compared to non-pregnant women (8).

Serum vitamin B<sub>12</sub> is increased by more than 40% at 6 weeks postpartum and remains so as long as the mothers are lactating. An additional moderate increase in the metabolic markers plasmas tHcy and MMA is observed,

suggestive of maternal intracellular vitamin B<sub>12</sub> depletion, despite high serum concentrations (8). Vitamin B<sub>12</sub> concentrations in breast milk decrease during the lactational period with the highest levels seen in the first 4 weeks (45).

#### *Inborn errors of metabolism*

There are several inborn errors of vitamin B<sub>12</sub> metabolism, which can be roughly divided into genetic mutations affecting absorption and transport, and mutations affecting intracellular vitamin B<sub>12</sub> processing. The former category includes Intrinsic factor deficiency and Immerslund-Gräsbeck syndrome, which inhibits absorption either through the lack of intrinsic factor or a defect in the cobalamin receptor facilitating intestinal uptake, ultimately resulting in vitamin B<sub>12</sub> deficiency. Haptocorrin deficiency is considered benign and results in low serum vitamin B<sub>12</sub> and normal concentrations of the metabolic markers. The inborn errors affecting intracellular processing are categorized based on whether they result in accumulation of Hcy and/or MMA. CblF, CblJ, CblC, CblX, or CblD are characterized by increased levels of both tHcy and MMA, and CblE and CblG yield isolated hyperhomocysteinemia, whereas CblA and CblB yield isolated methylmalonic aciduria. Most inborn errors of vitamin B<sub>12</sub> metabolism are rare, and the most common mutation is CblC, which has been reported in >500 patients (46).

#### **Assessment of nutrient status**

Vitamin B<sub>12</sub> status is judged based on the estimated dietary vitamin B<sub>12</sub> intake and relevant symptoms that may suggest vitamin B<sub>12</sub> deficiency or insufficiency. Biomarkers of vitamin B<sub>12</sub> status include serum vitamin B<sub>12</sub> and holoTC (bioavailable fraction in the circulation), while the functional biomarkers, plasma total tHcy and MMA, reflect intracellular vitamin B<sub>12</sub> function. Because B<sub>12</sub> is essential for folate metabolism, it is also important to consider folate status.

Many laboratories still use the 2.5 percentile, ranging from 145 to 200 pmol/L for serum vitamin B<sub>12</sub>, to define B<sub>12</sub> deficiency. A reference interval is typically defined as the 95% interval between the lower 2.5th and the upper 97.5th percentiles derived from the distribution of results from an apparently healthy reference population (47). Reference intervals merely describe the vitamin B<sub>12</sub> status in a specific population and will differ according to the diet in the tested population. One needs clinical decision to define deficiency.

Clinical decision limits define a value above or below a threshold associated with a significantly higher risk of adverse clinical outcomes or diagnostic for the presence of a specific disease (47). Vitamin B<sub>12</sub> deficiency associated with classic hematological and neurological

manifestations is relatively uncommon, and as clinical symptoms of vitamin B<sub>12</sub> deficiency generally are subtle and difficult to diagnose, a diagnostic delay of several months is common among infants, children, and adults (2, 5), something which is particularly unfortunate in young infants.

Various algorithms have been proposed in order to separate vitamin B<sub>12</sub> insufficiency from deficiency (2); however, on the cellular levels, there is a physiologic transition from inadequate to clearly deficient states of vitamin B<sub>12</sub>. In both children and adults, the metabolic markers tHcy and MMA start to increase when serum vitamin B<sub>12</sub> falls below ~500–550 pmol/L, indicating suboptimal intracellular vitamin B<sub>12</sub> stores, with a steeper increase in both markers when serum vitamin B<sub>12</sub> falls below 250–275 pmol/L, indicating biochemical deficiency (48, 49). Studies in humans show that biomarkers for DNA damage, such as hypomethylation, chromosome breaks, uracil incorporation, and micronucleus formation, are minimized when serum concentration of vitamin B<sub>12</sub> is greater than 300 pmol/L (50), indicating that also vitamin B<sub>12</sub> insufficiency is harmful and should be treated. It is obviously not good medical practice to wait until the store of an essential micronutrient is depleted.

In the following, we will comment on the individual biomarkers and the biological variations that should be taken into account when employing them. All four vitamin B<sub>12</sub> biomarkers have limitations as standalone markers, and there is currently no clear agreement on the decision limits for any of these parameters concerning vitamin B<sub>12</sub> deficiency or insufficiency (1, 2). Current diagnostic practice may be too complicated and focused on separating insufficiency from deficiency. As vitamin B<sub>12</sub> is an essential micronutrient it is also important to focus on what is an adequate status.

#### *Serum or plasma cobalamin*

Serum vitamin B<sub>12</sub> is still the primary marker of vitamin B<sub>12</sub> status among children, adults, and pregnant women (2). The assays used to analyze human samples are widely available, cheaper than all the other vitamin B<sub>12</sub> markers, and specific for biologically active cobalamins. A limitation of this biomarker is that it measures the total circulating cobalamin, of which the majority (70–90%) is bound to haptocorrin, which may be affected by high estrogen levels, cancer, and genetically low haptocorrin levels. In these circumstances, serum vitamin B<sub>12</sub> concentrations may not accurately reflect intracellular vitamin B<sub>12</sub> status (1, 2).

The serum concentrations are not substantially affected by recent dietary intake, meaning prandial status is not a major concern when collecting samples. In the longer term (months), the concentrations increase with increasing



habitual dietary intake and have been reported to plateau at vitamin B<sub>12</sub> intakes of 7–10 µg (14). A meta-analysis of 37 randomized controlled trials (RCTs) and 19 observational studies aimed to quantify the dose-response relationship and reported that each doubling of vitamin B<sub>12</sub> intake increased circulating concentrations by 11% (95% CI 9.4–12.5%). However, substantial heterogeneity between the studies, not explained by study design, age, or vitamin B<sub>12</sub> dose, warrants caution when interpreting the reported associations (15).

Diagnostic cutoffs vary from below 148 up to 300 pmol/L depending on the outcome (2). The lower decision levels (<148 pmol/L) are generally based on the presence of clinical deficiency symptoms, while studies show that DNA damage is minimized when serum vitamin B<sub>12</sub> is >300 pmol/L (50). A steep increase in both plasma tHcy and MMA occurs when serum vitamin B<sub>12</sub> falls below 250–275 pmol/L, indicating biochemical insufficiency and a cellular need for vitamin B<sub>12</sub>.

All vitamin B<sub>12</sub> markers are decreased during pregnancy, so specific decision limits must be used. A maternal vitamin B<sub>12</sub> concentration >275 pmol/L (measured by immunoassay) in week 18 of pregnancy has been recommended to secure an optimal infant vitamin B<sub>12</sub> status for the first 6 months of life (8, 51).

#### HoloTC

Circulating holoTC represents the active fraction (10–30%) of serum vitamin B<sub>12</sub>, and the test has been shown to provide a somewhat better or an equal discrimination compared to serum vitamin B<sub>12</sub>, also depending upon the decision limits used (52). Unlike the other biomarkers, HoloTC increases in the postprandial period and is therefore a better indicator of recent intake (14). This is taken advantage of when evaluating vitamin B<sub>12</sub> absorption with the CobaSorb test, measuring the holoTC response after an oral cyanocobalamin dose (53). Circulating concentrations also increase with increasing habitual vitamin B<sub>12</sub> intakes and stabilizes at intakes >4 µg (14, 54).

HoloTC is reported to be affected by genetic polymorphisms (55) and renal function (56). There is a lack of consensus concerning decision limits with current suggested limits for deficiency in adults <35–40 pmol/L, based on the relation between holoTC and serum MMA concentration (14). While the other biomarkers of vitamin B<sub>12</sub> status change during pregnancy, holoTC has been reported to be unaffected (14). However, plasma holoTC concentrations are reported to be reduced by 24% from preconception to pregnancy week 20, with an additional 17% reduction at labor (57). As long as holoTC decision limits for pregnant women and children are missing, serum vitamin B<sub>12</sub> remains the preferred biomarker in these groups.

#### Plasma MMA

Plasma MMA also accumulates with insufficient intracellular vitamin B<sub>12</sub> status, due to reduced activity of methylmalonyl-CoA mutase. As this reaction is not directly influenced by other micronutrients, MMA has, for a long time, been considered the most specific and informative of the vitamin B<sub>12</sub> biomarkers. However, MMA concentrations are influenced by several other factors. Less than 17% of the MMA concentrations in blood are determined by cobalamin, renal function, age, and gender, and MMA is therefore not the perfect vitamin B<sub>12</sub> marker (49). Currently, higher concentrations are seen with impaired renal function and with increasing age, especially after 70 years (49). The latter is not entirely explained by age-related decline in kidney function. Furthermore, two genetic variants have been reported to explain 12% of the variance in circulating MMA (58). In a meta-analysis, a 7% decrease (95% CI 4–10%) in MMA was reported for each doubling in vitamin B<sub>12</sub> intake (15).

In adults with normal kidney functions, clinical decision limits in the range >0.376 to >0.271 µmol/L have been suggested as markers for vitamin B<sub>12</sub> deficiency. In order to rule out impaired renal function as the cause of elevated MMA, renal function should be evaluated (14).

MMA has been reported to increase throughout pregnancy, indicating vitamin B<sub>12</sub> insufficiency (8, 59). However, compared to non-pregnant women, lower concentrations have been reported in pregnant women (8). In infants and toddlers, the MMA concentrations are higher across the entire vitamin B<sub>12</sub> spectrum compared to older children and adults (43), in this age-group tHcy should be used as a vitamin B<sub>12</sub> marker. In older children age-specific decision limits should be used.

#### Plasma tHcy

Plasma tHcy accumulates with insufficient intracellular vitamin B<sub>12</sub> status, due to reduced function of methionine synthase, referred to as hyperhomocysteinemia. Plasma tHcy concentrations are not a specific vitamin B<sub>12</sub> status marker, as it also increases with insufficient intakes of folate and to a lesser degree with low intake of vitamin B6.

Plasma tHcy concentrations increase with age and decreased renal function and are influenced by sex and pregnancy, so it is necessary to use age-specific decision limits (2). The most common decision limit to define hyperhomocysteinemia is >15 µmol/L in adults.

Plasma tHcy is the preferable metabolic marker for vitamin B<sub>12</sub> status in infants and toddlers, whereas in older children and adults, tHcy is primarily a folate marker, and MMA is the preferable vitamin B<sub>12</sub> marker (2). In infants and toddlers up to 3 years, a plasma tHcy concentration >6.5 µmol/L indicates vitamin B<sub>12</sub> deficiency (41, 44).

### Dietary intake in Nordic and Baltic countries

Vitamin B<sub>12</sub> only occurs in foods of animal origin and in fortified foods. Meat, liver, eggs, dairy products, fish, and shellfish are particularly good sources and are the main vitamin B<sub>12</sub> sources in the average diet.

The average vitamin B<sub>12</sub> intake in the Nordic countries, according to national dietary surveys, ranged from 4.0–6.4 to 5.5–8.9 µg per day in adult women and men, respectively (60).

#### Vegans/Vegetarians

Vegetarian and, especially, vegan diets tend to contain low or no amounts of vitamin B<sub>12</sub>, and these diets are associated with an increased risk of vitamin B<sub>12</sub> deficiency unless adequate vitamin supplementation is implemented (23, 61–64). Many vegan organizations have set detailed instructions for taking a supplement, and the main principle is that the less often the vitamin B<sub>12</sub> supplement is taken, the greater the dose must be; non-prescription vitamin B<sub>12</sub> supplements containing up to 1,000 µg vitamin B<sub>12</sub> per tablet are currently available in pharmacies.

Plant foods might contain trace amounts due to bacterial or soil contamination or as a result of fermentation. The amounts of vitamin B<sub>12</sub> in seaweeds cannot be considered a reliable or relevant source of vitamin B<sub>12</sub> for humans (65). Some plant-based milk substitutes are fortified with vitamin B<sub>12</sub> and can be an important source of vitamin B<sub>12</sub> in vegans. The nutritional quality of plant-based meat analogues on the Swedish market was assessed and published in 2022. The authors showed that vitamin B<sub>12</sub> contents in fortified products differed and were lower, similar, or higher compared to meat references (66).

A recent study of vitamin B<sub>12</sub> status in Norwegian vegans and vegetarians showed that average daily supplement use and older age were predictors of higher serum B<sub>12</sub> concentrations. In this study, none of the participants were considered to be B<sub>12</sub> depleted; however, low serum B<sub>12</sub> concentration was found in 14% of the participants (67).

#### Pregnant women

In pregnancy week 18, women from the Norwegian Mother, Father and Child Cohort Study (*MoBa*) had a median vitamin B<sub>12</sub> intake of 8.5 µg/day from diet and supplements (10).

#### Vitamin B<sub>12</sub> in breast milk and formula and intake in infants

In breast milk from Danish mothers, most of them took daily multivitamin supplements containing 1.0–4.5 µg cobalamin, the median (25<sup>th</sup> to 75<sup>th</sup> percentile) vitamin B<sub>12</sub> concentration at 2 weeks was 0.76 (0.21–1.88) nmol/L, at 4 months was 0.29 (0.14–0.69) nmol/L, and at 9 months was 0.44 (0.16–1.94) nmol/L (45). In breast milk from Norwegian mothers, users of supplements versus non-users, the median (25<sup>th</sup> to 75<sup>th</sup> percentile) vitamin B<sub>12</sub> concentration at 6 weeks was 0.34 (0.23–0.46) versus 0.29 (0.16–0.36) nmol/L, at 4 months was 0.31 (0.21–0.40) versus 0.27 (0.18–0.48) nmol/L, and at 6 months was 0.35 (0.22–0.54) versus 0.24, (0.19–0.47) nmol/L. Mean vitamin B<sub>12</sub> content in 11 different formula milk was 2.2 µg/L in a Danish study published in 2016 (68). Table 1 shows the estimated vitamin B<sub>12</sub> intake in infants from breast milk from these Danish and Norwegian mothers and from formula milk. Based on these data, formula-fed infants receive from 2.1 (at 2 weeks) up to 5.2 (at 4 months) times more vitamin B<sub>12</sub> per day compared to breast-fed infants.

Despite a higher milk intake with increasing age, daily vitamin B<sub>12</sub> intake is reduced from 2 weeks to 6 months in exclusively breast-fed infants (8, 45, 69, 70).

In a Norwegian study, the estimated vitamin B<sub>12</sub> intake at 12 months was median 1.7 (IQR 1.1–2.2) µg per day from solid food and breastmilk versus 2.3 (1.8–3.0) µg per day from solid food and formula (70).

#### Infants born premature or with LBW

Infants born premature or with a low birthweight have smaller vitamin B<sub>12</sub> stores and an increased risk of deficiency during the first year of life (71). However, as many premature and LBW infants have

**Table 1.** Median (IQR) vitamin B<sub>12</sub> intake from breast milk versus formula milk in infants

Parameters	2 weeks	6 weeks	4 months	6 months	9 months
Mean weight, kg	3.7	4.8	6.7	7.6	8.6
Mean milk intake/day	150 ml/kg	130 ml/kg	120 ml/kg	120 ml/kg	100 ml/kg
Median infant vitamin B <sub>12</sub> intake from:					
Danish mothers, +Suppl. (45)	0.57 µg/day		0.31 µg/day		0.51 µg/day
Norwegian mothers (8)		0.29 µg/day	0.34 µg/day	0.43 µg/day	
+Suppl.		0.24 µg/day	0.29 µg/day	0.30 µg/day	
–Suppl.					
Formula (68)	1.22 µg/day	1.37 µg/day	1.77 µg/day	2.00 µg/day	1.89 µg/day

problems breastfeeding, this group is often mainly formula fed, something which will increase their vitamin B<sub>12</sub> intake and improve their status (7). In exclusively breast-fed premature or low birthweight infants, one must be aware of development of vitamin deficiency.

#### *Elderly*

Dietary surveys in the Nordic countries suggest that vitamin B<sub>12</sub> intake in the elderly is slightly higher compared to younger adults, with estimated intakes ranging from 5.2–6.6 to 5.2–10.8 µg/day in women and men, respectively. However, with increasing age, the production of both hydrochloric acid and intrinsic factor declines, putting elderly at increased risk of developing vitamin B<sub>12</sub> deficiency or insufficiency (2). It has been estimated that 10–30% of elderly have some form of vitamin B<sub>12</sub> malabsorption due to atrophic gastritis, which is often undiagnosed.

### **Health outcomes relevant for Nordic and Baltic countries**

#### *Deficiency*

There is currently no clear agreement on what symptoms constitute clinical vitamin B<sub>12</sub> deficiency and what decision limits for vitamin B<sub>12</sub> status one should use (2), but one guiding principle is that if the symptoms are alleviated by treatment with cobalamin, then it is likely that there was a degree of vitamin B<sub>12</sub> insufficiency. An adequate supply of vitamin B<sub>12</sub> is essential for normal development, neurological function, and blood formation (2). Vitamin B<sub>12</sub> deficiency in adults may result in neurological and psychiatric symptoms and/or macrocytic and megaloblastic anemia (2, 72, 91). Lindenbaum et al. reported vitamin B<sub>12</sub> deficient patients with psychiatric symptoms, but without anemia or macrocytosis (72). The reason why some patients mainly present with megaloblastic anemia and others with neurologic symptoms remains unknown. Neurological manifestations, such as numbness and paraesthesia in the extremities; loss of position and vibratory sensation; difficulty walking; depression and irritability; diminished cognitive function; and psychosis may be observed in the absence of hematologic disease, (72, 73). Subclinical deficiency detected by metabolic markers i.e. increased tHcy and MMA homocysteine and MMA, which are predictors of developmental and degenerative diseases, is much more common in the general population (50). The observed biochemical insufficiency at B<sub>12</sub> levels below ~500–550 pmol/L raises the question whether such vitamin B<sub>12</sub> insufficiency has clinical consequences (48). A review has shown a high prevalence of vitamin B<sub>12</sub> insufficiency

in the world and has identified several examples of adverse clinical outcomes, mainly in relation to functions of the nervous system (74). Others have suggested that vitamin B<sub>12</sub> deficiency/insufficiency is related to increased prooxidant and reduced antioxidant status (32).

#### *Vitamin B<sub>12</sub> deficiency during pregnancy and infancy*

Vitamin B<sub>12</sub> deficiency in women of fertile age is associated with infertility, early pregnancy loss (75), increased prevalence of preeclampsia, preterm birth (76) possibly LBW (27, 29), and a moderate risk of neural tube defects (77). A review published including 23 studies from 1961 to 2017 reported a positive effect of vitamin B<sub>12</sub> on semen quality, including increasing sperm count, enhanced sperm motility, and reduced sperm DNA damage (78).

Maternal vitamin B<sub>12</sub> deficiency during pregnancy is associated with vitamin B<sub>12</sub> deficiency in the infant (8). Symptoms of vitamin B<sub>12</sub> deficiency in children differ with age, presenting a continuum from subtle developmental delay to life-threatening clinical conditions. The first months of life is the most rapid period of brain development, with a brain growth ranging per day from 1% per day and gradually decreasing to 0.4% at 3 months (79). An optimal micronutrient status is particularly important during this period. In infants, symptoms of vitamin B<sub>12</sub> deficiency include irritability, failure to thrive, apathy, anorexia, gastrointestinal reflux, constipation, and developmental delay and regression (5, 6), all of which respond remarkably and rapidly to supplementation (6). Symptoms of vitamin B<sub>12</sub> deficiency are difficult to detect in all age groups, but particularly in infants, and there tends to be a diagnostic delay up to 4 months in this age group (5, 9). Several reports show that even moderate deficiency in children may be harmful, and long-term consequences of neurological deterioration may persist after vitamin B<sub>12</sub> deficiency has been treated (9). Even in well-nourished children, a low vitamin B<sub>12</sub> concentration has been associated with a lower score for psychomotor development. A randomized controlled vitamin B<sub>12</sub> intervention study in infants with moderate B<sub>12</sub> demonstrated significantly improved motor development in only 3 weeks (6). In Danish children aged 36 months with a mean (SD) serum vitamin B<sub>12</sub> concentration of 653 (240) pmol/L, an increase of 100 pmol/l in serum vitamin B<sub>12</sub> corresponded with a 1.5 increase in total Ages and Stages Questionnaire (ASQ) score, a measure of psychomotor development, raising a question about what is the optimal serum vitamin B<sub>12</sub> concentration in young children. Children in the lowest vitamin B12 quartile had about 10 ASQ scores lower compared to the other quartiles (80).

#### *Toxicity*

As long as vitamin B<sub>12</sub> is taken orally, either in food or in supplements containing a moderate vitamin B<sub>12</sub> content

(<50 µg), the body will actively regulate the uptake in ileum, and excess vitamin B<sub>12</sub> will be excreted in urine. To patients with confirmed vitamin B<sub>12</sub> deficiency due to absorption problems, a peroral dose of 4,000 cyanocobalamin µg daily for 1 month, followed by 1–2,000 µg per day, is now recommended as an alternative to vitamin B<sub>12</sub> injection therapy. There are intake known severe adverse effects of excess vitamin B<sub>12</sub> (12). Potential negative effects of megadoses, i.e. on the microbiome, are largely unknown, but one study reported that vitamin B<sub>12</sub> supplementation may give acne development due to changes in the skin microbiome (81).

### Obesity

The association between circulating vitamin B<sub>12</sub> concentrations and the body mass index (BMI) was investigated in a systematic review of 19 observational studies including 7,055 participants (33). Overall, it was reported that those with BMI categorized as overweight or obese had mean 21–56 pmol/L lower serum vitamin B<sub>12</sub> concentrations compared to those with normal weight. In meta-regression analyses, no inverse or J-shaped association was established, but the authors concluded that the results from the direct pairwise comparisons supported further investigation on the association between vitamin B<sub>12</sub> status and overweight/obesity. Lower vitamin B<sub>12</sub> levels with higher BMI are also reported in pregnant women (82).

### Cardiovascular disease (CVD) and diabetes type 2

In a Cochrane review of 15 RCTs, Marti-Carvajal et al. reported on the effect of homocysteine-lowering intervention with B-vitamin supplements (vitamin B<sub>6</sub>, folate, and vitamin B<sub>12</sub> alone or in combination) on cardiovascular disease outcomes and mortality (20). Compared to placebo, they reported reduced risk of stroke (RR [95% CI] 0.90 [0.82–0.99]), but no effect on myocardial infarction risk (1.02 [0.95–1.10]) or all-cause mortality (1.01 [0.96–1.06]). Rafnsson et al. performed a systematic review of 7 cohort studies found only very limited evidence for the associations between low serum vitamin B<sub>12</sub> and increased risk of mortality and morbidity from either cardiovascular diseases or type 2 diabetes in adults (26).

### Cancer

Observational studies have found an association between higher circulating vitamin B<sub>12</sub> concentrations and increased risk of prostate (83) and total cancers (84), and a high serum vitamin B<sub>12</sub> has been proposed as a cancer biomarker (85). The elevated vitamin B<sub>12</sub> levels appear to be due to increased haptocorrin concentrations and not to raised holoTC (83, 86). Furthermore, clinical trials do not support an association between higher vitamin B<sub>12</sub> intakes and cancer risk (87–89). A meta-analysis

of 18 RCTs including approximately 75,000 individuals found that supplements containing B vitamins, including 20 to 2,000 µg/day vitamin B<sub>12</sub>, had little or no effect on cancer incidence, cancer deaths, or all-cause mortality (88). A dose-response meta-analysis including data from 10,600 patients suggested an inverse association between both total and dietary vitamin B<sub>12</sub> intakes and colorectal cancer risk (90). This was supported in a meta-analysis of 13 case-control studies, reporting an inverse association between circulating vitamin B<sub>12</sub> and colorectal cancer risk (not statistically significant), and a positive association with circulating tHcy (30). However, a meta-analysis of 10 observational studies with 3,164 cases observed a higher risk of esophageal cancer when comparing the highest vs lowest intake (OR [95% CI] 1.30 [1.05–1.62]), and a linear dose-response per 1 µg/d increment (1.02 [1.00–1.03]) (25).

### Osteoporosis and bone health

The association of vitamin B<sub>12</sub> status with bone mineral density and osteoporotic fractures was investigated in a systematic review by Macedo et al., including 6 longitudinal studies, 9 cross-sectional studies, and two RCTs (19). The authors concluded that the potential effect of vitamin B<sub>12</sub> deficiency on bone health and the underlying mechanisms warrant further research, especially in vulnerable groups such as postmenopausal and elderly women.

### Mental health

Whereas most studies question the effect of vitamin B<sub>12</sub> on mental health in the elderly, positive associations are reported in children (80). Data on vitamin B<sub>12</sub> status and associated cognitive function and depression in elderly people are conflicting. A Cochrane review from 2003 found that the evidence for any effect of vitamin B<sub>12</sub> on improving the cognitive function of people with dementia and low serum vitamin B<sub>12</sub> levels was insufficient (92). The same conclusion was reached in a systematic review and meta-analysis of 19 RCTs (16), concluding that supplementation of cobalamin, vitamin B<sub>6</sub>, or folic acid alone, or in combination, did not appear to improve cognitive functions in individuals with or without cognitive impairment at baseline. The same authors performed an updated systematic review of 31 RCTs, and in a meta-analysis of 12 RCTs using the Mini Mental State Examination, they observed no meaningful effect in individuals with, and no effect in individuals without, cognitive impairment at baseline (31). Accordingly, there is limited evidence from observational studies to suggest an association of vitamin B<sub>12</sub> status with cognitive decline or dementia in elderly people (21, 93). In a systematic review of observational studies, the authors concluded that there was insufficient evidence to establish an overall association, but it was highlighted that all four



studies that used more specific biomarkers of vitamin B<sub>12</sub> status (MMA and holoTC) showed consistent associations between poor vitamin B<sub>12</sub> status and increased risk of cognitive decline or dementia diagnosis (22). Another systematic review identified poor vitamin B<sub>12</sub> status as being associated with increased risk of vascular dementia, primarily based on cross-sectional studies (24). Treatment with a combination of folic acid and vitamin B<sub>12</sub> has also shown cognitive benefits in people with mild cognitive impairment (94, 95).

Vitamin B<sub>12</sub> insufficiency has been found to increase the risk of depression in the elderly (96), and a recent review concluded that early vitamin B<sub>12</sub> supplementation can delay the onset of depression and improve the effect of anti-depressants (97). On the other hand, a systematic review of 11 RCTs concluded that short-term treatment with folic acid and/or vitamin B<sub>12</sub> did not improve depressive symptoms. However, the results suggested that longer-term treatment may decrease the risk of relapse or onset of symptoms in those at increased risk (98). A systematic review pooling data from 10 observational studies reported consistently lower circulating vitamin B<sub>12</sub> concentrations in patients with Parkinson's disease (overall standardized mean difference [95% CI] -0.38 [-0.51, -0.25]) (28). However, no association was observed for dietary vitamin B<sub>12</sub> intake and Parkinson's disease risk.

#### Eye disease

In a systematic review and meta-analysis of 11 case-control studies, including 1,072 cases, Huang et al. reported that tHcy was higher (mean difference [95% CI] 2.67 [1.6–3.74] µmol/L), and serum vitamin B<sub>12</sub> was lower (64.2 [19.3–109] pmol/L) in age-related macular degeneration (AMD) cases compared to controls (17). Another systematic review and meta-analysis by Li et al. included 9 studies reporting on the serum vitamin B<sub>12</sub> levels and different types of glaucoma (18). In random-effect meta-analyses, they reported no statistically significant differences in serum vitamin B<sub>12</sub> between cases and controls.

#### Requirement and RIs

The RI of vitamin B<sub>12</sub> published in NNR 2004 and 2012 was largely based on a study published in 1958, where 20 patients with pernicious anemia were treated with an intramuscular dose of 0.5–2.0 µg vitamin B<sub>12</sub> per day, and the outcome parameter was normalizing of hematological status (99). An average physiological requirement of vitamin B<sub>12</sub> was set at 0.7 µg/day, based on this and other studies (38, 100). With correction for absorption efficiency (50%), the average requirement (AR) was set at 1.4 µg/day for adults. By assuming a CV of 15% and adding two SD to allow for individual variation, the RI for adults was set at 2 µg/day. The same recommendations applied

to elderly and pregnant women, while an additional 0.6 µg/day was recommended to lactating women. The recommendations for children were based on an assumed requirement of 0.05 µg/kg body weight/day, and the RIs were set at 0.8 (2–5y), 1.3 (6–9y), and 2.0 (10–13y) µg/day. The NNR2012 RI was unchanged from the recommendations in NNR2004.

The NNR2012 DRVs were comparable to most other vitamin B<sub>12</sub> intake recommendations by authoritative scientific bodies worldwide (14), with one notable exception. In 2015, the EFSA Panel concluded that it is impossible to determine an AR and RI and, therefore, set an adequate intake (AI) for vitamin B<sub>12</sub> at 4 µg/day for men and women >14y, based on available data on different vitamin B<sub>12</sub> biomarkers (101). An additional 0.5 µg/d was added for pregnancy (AI 4.5 µg/day), and 1 µg/day was added during lactation (AI 5.0 µg/day) based on fetal accumulation of vitamin B<sub>12</sub> and excretion through breast milk. For children, the AIs were extrapolated from adult values and were set at 1.5 µg/day from 7 months of age.

The *de novo* NNR2023 systematic review concluded that for all included populations (pregnancy, lactation, infants, elderly, and vegetarians and vegans), there was insufficient evidence to assess whether habitual intakes in line with the previously RIs (NNR2012) were sufficient to maintain adequate status (35). No eligible studies were identified for young adults or children other than breast-fed infants.

#### Adults

Daily vitamin B<sub>12</sub> losses in apparently healthy adults and elderly probably range from 1.4 to 5.1 µg. A systematic review estimated that the vitamin B<sub>12</sub> intakes needed to compensate for these losses ranged from 3.8 to 20.7 µg (11). However, these estimates should be interpreted with caution, as the rate-limited absorption from a single dose was not considered. To better estimate the dose needed, both the bioavailability and the division of the dose throughout the day must be considered. Based on biomarker concentrations, it has been reported that in healthy young adults and postmenopausal women, circulating concentrations of cobalamin, homocysteine, and MMA appears to plateau at intakes of 4–10 µg/day (54, 102, 103). This suggests that previous RIs (NNR2012) may be insufficient to maintain optimal biomarker status. Based on available data on different biomarkers of vitamin B<sub>12</sub> status, a vitamin B<sub>12</sub> intake in the range of 5–7 µg/day seems to be adequate for adults; however, no consensus exists regarding what should be considered optimal in terms of metabolic profile.

#### Elderly

In NNR2012, the RI for elderly was the same as for younger adults. This is in line with the Institute of

Medicine (IOM), who emphasize that because dietary vitamin B<sub>12</sub> can be assumed to be poorly absorbed due to atrophic gastritis, increasing the recommendations was unlikely to be sufficient.

#### *Vegetarian and vegan diets*

As vitamin B<sub>12</sub> only occurs in foods of animal origin, vegan diets will eventually result in vitamin B<sub>12</sub> deficiency unless supplemented. However, lower intake and status are also observed with less restrictive diets limiting animal food (104). It should be emphasized that symptoms of deficiency may not occur until years after adopting such diets. As repleting the body stores through supplements would take a long time, initial treatment with injections is warranted once symptoms of deficiency are present.

#### *Pregnancy and lactation*

A maternal vitamin B<sub>12</sub> concentration >275 pmol/L in week 18 of pregnancy has been recommended to ensure an adequate infant vitamin B<sub>12</sub> status for the first 6 months of life (8, 49). Additionally, serum vitamin B<sub>12</sub> concentration above 300 pmol/L in early pregnancy has been associated with lower incidence of neural tube defects (105).

In pregnancy week 18, women from the Norwegian MoBa study reported a median vitamin B<sub>12</sub> intake of 8.5 µg/day from diet and supplements (10). Median (IQR) serum vitamin B<sub>12</sub> concentration in 2,911 women from the MoBa study was 309 (249–378) pmol/L (106). This study may suggest that a vitamin B<sub>12</sub> intake of 8.5 µg/d can be expected to give more than 50% of the women a sufficient vitamin B<sub>12</sub> status in pregnancy week 18.

#### *Infants*

##### **Infants born at term with an appropriate weight for gestational age**

The calculated AI for infants aged 0–6 months ranges from 0.3 to 0.5 µg/day and is based on the assumption that breast milk contains sufficient vitamin B<sub>12</sub> for optimal health during this period of life. The data for estimated AI include an average intake of breast milk (~800 ml/d) and an average vitamin B<sub>12</sub> concentration in breast milk (~0.45 µg/L).

Only 28% of infants at 6 weeks and 34% at 4 months achieve the RI of 0.4 µg per day from breast milk (8). Additionally, whether this amount of vitamin B<sub>12</sub> is enough will also depend on the infant vitamin B<sub>12</sub> stores, which will be low in infants born to women with vitamin B<sub>12</sub> deficiency due to gastrointestinal disease or a diet with a low content of animal food. Formula-fed infants receive from 1.2 to 2 µg vitamin B<sub>12</sub> per day from 2 weeks to 6 months (Table 1) and are reported to have lower plasma tHcy concentrations, indicating a better vitamin B<sub>12</sub> status compared to exclusively breast-fed infants (9).

For infants aged 7–12 months, the recommended AI ranges from 0.7 to 1.5 µg/day. EFSA has the highest AI (1.5 µg/day), based on an extrapolation from the AI for adults using allometric scaling and applying a growth factor (93). At 12 months, infants with median vitamin B<sub>12</sub> intake of 2.3 (25<sup>th</sup>–75<sup>th</sup> percentile 1.8–3.0) µg per day had a lower geometric mean plasma tHcy: 4.5 (95%CI 4.1–4.9) µmol/L, compared to infants with a lower median vitamin B<sub>12</sub> intake (1.7 (1.1–2.2) µg per day) plasma tHcy 5.5 (5.1–5.8) µmol/L. This study suggests that a vitamin B<sub>12</sub> intake of approximately 2.3 µg per day should be sufficient for most infants aged 12 months.

##### **Infants born premature or with LBW (<2,500 g)**

Infants born premature or with a LBW have an increased risk of vitamin B<sub>12</sub> deficiency during the first year of life (71). One study showed that breast-fed and formula-fed LBW infants had similar plasma tHcy until 20 postnatal days; after this, plasma tHcy subsequently increased in breast-fed infants, but not in formula-fed infants (107). The vitamin B<sub>12</sub> intake would be 0.17 to 0.23 µg vitamin B<sub>12</sub> per day (based on a vitamin B<sub>12</sub> concentration in breast milk of ~0.45 µg/L and 0.4–0.5 L milk intake/day) in the breast-fed infant. A formula-fed infant would get 0.8 to 1 µg vitamin B<sub>12</sub> per day (based on 2.2 µg/L vitamin B<sub>12</sub> and 0.4–0.5 L milk intake/day) during the period from 3 to 6 weeks. As plasma tHcy did not increase in the formula-fed infants, 0.8 to 1.0 µg vitamin B<sub>12</sub> per day seems to be the necessary amount of vitamin B<sub>12</sub> intake for LBW infants aged 3 to 6 weeks.

Among Norwegian infants with a birth weight between 2 and 3 kg, those who were exclusively breast-fed for >1 month had lower vitamin B<sub>12</sub> and higher tHcy levels at 4 and 6 months compared to infants who were fed formula. Additionally, the breast-fed infants had lower gross motor scores at 6 months compared to the formula-fed infants (7), indicating that even moderate vitamin B<sub>12</sub> insufficiency may impair neurodevelopment. Given a vitamin B<sub>12</sub> concentration in formula of 2.2 µg/L and a milk intake of 0.120 L/kg/day, the estimated vitamin B<sub>12</sub> intake associated with a better metabolic status and neurodevelopment would range from 1.1 µg/day at 4 months to 1.5 µg/day at 6 months.

#### *Children*

In healthy, non-breast-fed Norwegian toddlers at age 24 months, a plateau in serum vitamin B<sub>12</sub> and holoTC was reached at a vitamin B<sub>12</sub> intake of ~3 µg/day (108). Neither MMA nor tHcy concentrations decreased with increasing vitamin B<sub>12</sub> intakes, which may indicate that a vitamin B<sub>12</sub> intake of ~3 µg/day at this age secures an adequate vitamin B<sub>12</sub> status.

In a survey published in 2015, Ungkost 3 (109), vitamin B<sub>12</sub> intake in Norwegian children aged 9 years was mean 4.9 (SD 2.2) µg/day, and that of aged 13 years was mean 5.3 (SD 3.7) µg/day (Ungkost 3, 2015). In a study published in 2003 (110), Norwegian children aged 1–10 years had a median serum vitamin B<sub>12</sub> of 551 (25th–75th percentile 456, 683) pmol/L, for the age group 10.5–15 years, it was 436 (295, 529) pmol/L, and for the age group 15.5–19 years, it was 369 (294, 452) pmol/L. As a serum vitamin B<sub>12</sub> >300 pmol/L indicates an adequate vitamin B<sub>12</sub> status, approximately 25% of the older children (>10.5 years) had a risk of vitamin B<sub>12</sub> deficiency, whereas this was not an issue for the younger children. In Canadian children aged 6 years and adolescents, a daily consumption of vitamin B<sub>12</sub> supplements was associated with higher serum vitamin B<sub>12</sub> concentrations, with no additional increase in serum vitamin B<sub>12</sub> at doses above 10 µg/day (111).

Based on these studies, a vitamin B<sub>12</sub> intake from 3 to 4.9 µg/day may be adequate for younger children, whereas children >10 years may need a vitamin B<sub>12</sub> intake in the range of 5.3–10 µg/day.

### Conflict of interest and funding

The authors have not received any funding or benefits from industry or elsewhere to conduct this study.

### References

- Hannibal L, Lysne V, Bjørke-Monsen AL, Behringer S, Grünert SC, Spiekerkoetter U, et al. Biomarkers and algorithms for the diagnosis of vitamin B12 deficiency. *Front Mol Biosci* 2016; 3: 27. doi: 10.3389/fmolb.2016.00027
- Green R, Allen LH, Bjørke-Monsen AL, Brito A, Gueant JL, Miller JW, et al. Vitamin B12 deficiency. *Nat Rev Dis Prim* 2017; 3: 17040. doi: 10.1038/nrdp.2017.40
- Allès B, Baudry J, Méjean C, Touvier M, Péneau S, Hercberg S, et al. Comparison of Sociodemographic and Nutritional Characteristics between Self-Reported Vegetarians, Vegans, and Meat-Eaters from the NutriNet-Santé Study. *Nutrients*. 2017; 9(9): 1023. doi: 10.3390/nu9091023. PMID: 28926931; PMCID: PMC5622783.
- Sanne, I., Bjørke-Monsen, AL. Lack of nutritional knowledge among Norwegian medical students concerning vegetarian diets. *J Public Health (Berl.)* 2022; 30: 495–501. doi: 10.1007/s10389-020-01327-7
- Dror DK, Allen LH. Effect of vitamin B12 deficiency on neurodevelopment in infants: current knowledge and possible mechanisms. *Nutr Rev* 2008; 66(5): 250–5. doi: 10.1111/j.1753-4887.2008.00031.x
- Torsvik I, Ueland PM, Markestad T, Bjørke-Monsen AL. Cobalamin supplementation improves motor development and regurgitations in infants: results from a randomized intervention study. *Am J Clin Nutr* 2013; 98(5): 1233–40. doi: 10.3945/ajcn.113.061549
- Torsvik IK, Ueland PM, Markestad T, Midttun O, Monsen AL. Motor development related to duration of exclusive breastfeeding, B vitamin status and B12 supplementation in infants with a birth weight between 2000–3000 g, results from a randomized intervention trial. *BMC Pediatr* 2015; 15(1): 218. doi: 10.1186/s12887-015-0533-2
- Varsi K, Ueland PM, Torsvik IK, Bjørke-Monsen AL. Maternal serum cobalamin at 18 weeks of pregnancy predicts infant cobalamin status at 6 months – a prospective, observational study. *J Nutr* 2018; 148(5): 738–45. doi: 10.1093/jn/nxy028
- Bjørke-Monsen AL, Ueland PM. Cobalamin status in children. *J Inher Metab Dis* 2011; 34(1): 111–19. doi: 10.1007/s10545-010-9119-1
- Haugen M, Brantsaeter AL, Alexander J, Meltzer HM. Dietary supplements contribute substantially to the total nutrient intake in pregnant Norwegian women. *Ann Nutr Metab* 2008; 52(4): 272–80. doi: 10.1159/000146274
- Doets EL, In 't Veld PH, Szczecinska A, Dhonukshe-Rutten RA, Cavelaars AE, van't Veer P, et al. Systematic review on daily vitamin B12 losses and bioavailability for deriving recommendations on vitamin B<sub>12</sub> intake with the factorial approach. *Ann Nutr Metab* 2013; 62(4): 311–22. doi: 10.1159/000346968
- SCF. Scientific Committee on Food. Opinion of the Scientific Committee on Food on the Tolerable Upper Intake Level of Vitamin B<sub>12</sub> 2000. [https://ec.europa.eu/food/sites/food/files/safety/docs/sci-com\\_scf\\_out80d\\_en.pdf](https://ec.europa.eu/food/sites/food/files/safety/docs/sci-com_scf_out80d_en.pdf). Accessed 04 March 2021.
- Blomhoff R, Andersen R, Arnesen EK, Christensen JJ, Eneroth H, Erkkola M, et al. *Nordic Nutrition Recommendations 2023*. Copenhagen: Nordic Council of Ministers; 2023.
- Allen LH, Miller JW, de Groot L, Rosenberg IH, Smith AD, Refsum H, et al. Biomarkers of nutrition for development (BOND): vitamin B-12 review. *J Nutr* 2018; 148(Suppl\_4): 1995S–2027S. doi: 10.1093/jn/nxy201
- Dullemeijer C, Souverein OW, Doets EL, van der Voet H, van Wijngaarden JP, de Boer WJ, et al. Systematic review with dose-response meta-analyses between vitamin B-12 intake and European micronutrient recommendations aligned's prioritized biomarkers of vitamin B-12 including randomized controlled trials and observational studies in adults and elderly persons. *Am J Clin Nutr* 2013; 97(2): 390–402. doi: 10.3945/ajcn.112.033951
- Ford AH, Almeida OP. Effect of homocysteine lowering treatment on cognitive function: a systematic review and meta-analysis of randomized controlled trials. *J Alzheimers Dis* 2012; 29(1): 133–49. doi: 10.3233/JAD-2012-111739
- Huang P, Wang F, Sah BK, Jiang J, Ni Z, Wang J, et al. Homocysteine and the risk of age-related macular degeneration: a systematic review and meta-analysis. *Sci Rep* 2015; 5: 10585. doi: 10.1038/srep10585
- Li S, Li D, Shao M, Cao W, Sun X. Lack of Association between Serum Vitamin B<sub>6</sub>, Vitamin B<sub>12</sub>, and Vitamin D Levels with Different Types of Glaucoma: A Systematic Review and Meta-Analysis. *Nutrients*. 2017; 9(6): 636. doi: 10.3390/nu9060636. PMID: 28635642; PMCID: PMC5490615.
- Macedo LLG, Carvalho C, Cavalcanti JC, Freitas B. Vitamin B12, bone mineral density and fracture risk in adults: a systematic review. *Rev Assoc Med Bras* 2017; 63(9): 801–9. doi: 10.1590/1806-9282.63.09.801
- Marti-Carvajal AJ, Sola I, Lathyris D, Dayer M. Homocysteine-lowering interventions for preventing cardiovascular events. *Cochrane Database Syst Rev* 2017; 8: CD006612. doi: 10.1002/14651858.CD006612.pub5
- Miles LM, Mills K, Clarke R, Dangour AD. Is there an association of vitamin B12 status with neurological function in older people? A systematic review. *Br J Nutr* 2015; 114(4): 503–8. doi: 10.1017/S0007114515002226

22. O'Leary F, Allman-Farinelli M, Samman S. Vitamin B(1)(2) status, cognitive decline and dementia: a systematic review of prospective cohort studies. *Br J Nutr* 2012; 108(11): 1948–61. doi: 10.1017/S0007114512004175
23. Obersby D, Chappell DC, Dunnett A, Tsiami AA. Plasma total homocysteine status of vegetarians compared with omnivores: a systematic review and meta-analysis. *Br J Nutr* 2013; 109(5): 785–94. doi: 10.1017/S000711451200520X
24. Perez L, Heim L, Sherzai A, Jaceldo-Siegl K, Sherzai A. Nutrition and vascular dementia. *J Nutr Health Aging* 2012; 16(4): 319–24. doi: 10.1007/s12603-012-0042-z
25. Qiang Y, Li Q, Xin Y, Fang X, Tian Y, Ma J, et al. Intake of Dietary One-Carbon Metabolism-Related B Vitamins and the Risk of Esophageal Cancer: A Dose-Response Meta-Analysis. *Nutrients*. 2018; 10(7): 835. doi: 10.3390/nu10070835. PMID: 29954131; PMCID: PMC6073467
26. Rafnsson SB, Saravanan P, Bhopal RS, Yajnik CS. Is a low blood level of vitamin B12 a cardiovascular and diabetes risk factor? A systematic review of cohort studies. *Eur J Nutr* 2011; 50(2): 97–106. doi: 10.1007/s00394-010-0119-6
27. Rogne T, Tielemans MJ, Chong MF, Yajnik CS, Krishnaveni GV, Poston L, et al. Associations of maternal vitamin B12 concentration in pregnancy with the risks of preterm birth and low birth weight: a systematic review and meta-analysis of individual participant data. *Am J Epidemiol* 2017; 185(3): 212–23. doi: 10.1093/aje/kww212
28. Shen L. Associations between B vitamins and Parkinson's disease. *Nutrients* 2015; 7(9): 7197–208. doi: 10.3390/nu7095333
29. Sukumar N, Rafnsson SB, Kandala NB, Bhopal R, Yajnik CS, Saravanan P. Prevalence of vitamin B-12 insufficiency during pregnancy and its effect on offspring birth weight: a systematic review and meta-analysis. *Am J Clin Nutr* 2016; 103(5): 1232–51. doi: 10.3945/ajcn.115.123083
30. Sun M, Sun M, Zhang L, Shi S. Colorectal polyp risk is linked to an elevated level of homocysteine. *Biosci Rep*. 2018; 38(2): 699. doi: 10.1042/BSR20171699. PMID: 29581245; PMCID: PMC5968185
31. Ford AH, Almeida OP. Effect of vitamin B supplementation on cognitive function in the elderly: a systematic review and meta-analysis. *Drugs Aging* 2019; 36(5): 419–34. doi: 10.1007/s40266-019-00649-w
32. van de Lagemaat EE, de Groot LCPGM, van den Heuvel EGHM. Vitamin B12 in Relation to Oxidative Stress: A Systematic Review. *Nutrients*. 2019; 11(2): 482. doi: 10.3390/nu11020482
33. Wiebe N, Field CJ, Tonelli M. A systematic review of the vitamin B12, folate and homocysteine triad across body mass index. *Obes Rev* 2018; 19(11): 1608–18. doi: 10.1111/obr.12724
34. Høyer A, Christensen JJ, Arnesen EK, Andersen R, Eneroth H, Erkkola M, et al. The Nordic Nutrition Recommendations 2022 - prioritisation of topics for de novo systematic reviews. *Food Nutr Res*. 2021; 65: 7828. doi: 10.29219/fnr.v65.7828
35. Bärebring L, Lamberg-Allardt C, Thorisdottir B, et al. Intake of vitamin B12 in relation to vitamin B12 status in groups susceptible to deficiency: a systematic review. *Food Nutr Res*. 2023; 67: 8626. doi: 10.29219/fnr.v67.8626
36. Wang H, Li L, Qin LL, Song Y, Vidal-Alaball J, Liu TH. Oral vitamin B12 versus intramuscular vitamin B12 for vitamin B12 deficiency. *Cochrane Database Syst Rev* 2018; 3: CD004655. doi: 10.1002/14651858.CD004655.pub3
37. Allen LH. Bioavailability of vitamin B12. *Int J Vitam Nutr Res* 2010; 80(4–5): 330–5. doi: 10.1024/0300-9831/a000041
38. Herbert V. Recommended dietary intakes (RDI) of vitamin B-12 in humans. *Am J Clin Nutr* 1987; 45: 671–8. doi: 10.1093/ajcn/45.4.671
39. Russell RM, Baik H, Kehayias JJ. Older men and women efficiently absorb vitamin B-12 from milk and fortified bread. *J Nutr* 2001; 131(2): 291–3. doi: 10.1093/jn/131.2.291
40. Chatthanawaree W. Biomarkers of cobalamin (vitamin B12) deficiency and its application. *J Nutr Health Aging* 2011; 15(3): 227–31. doi: 10.1007/s12603-010-0280-x
41. Bjørke Monsen AL, Ueland PM, Vollset SE, Guttormsen AB, Markestad T, Solheim E, et al. Determinants of cobalamin status in newborns. *Pediatrics* 2001; 108(3): 624–30. doi: 10.1542/peds.108.3.624
42. Minet JC, Bisse E, Aebischer CP, Beil A, Wieland H, Lutschg J. Assessment of vitamin B-12, folate, and vitamin B-6 status and relation to sulfur amino acid metabolism in neonates. *Am J Clin Nutr* 2000; 72(3): 751–7. doi: 10.1093/ajcn/72.3.751
43. Monsen AL, Refsum H, Markestad T, Ueland PM. Cobalamin status and its biochemical markers methylmalonic acid and homocysteine in different age groups from 4 days to 19 years. *Clin Chem* 2003; 49(12): 2067–75. doi: 10.1373/clinchem.2003.019869
44. Bjørke-Monsen AL, Torsvik I, Saetran H, Markestad T, Ueland PM. Common metabolic profile in infants indicating impaired cobalamin status responds to cobalamin supplementation. *Pediatrics* 2008; 122(1): 83–91. doi: 10.1542/peds.2007-2716
45. Greibe E, Lildballe DL, Streym S, Vestergaard P, Rejnmark L, Mosekilde L, et al. Cobalamin and haptocorrin in human milk and cobalamin-related variables in mother and child: a 9-mo longitudinal study. *Am J Clin Nutr* 2013; 98(2): 389–95. doi: 10.3945/ajcn.113.058479
46. Watkins D, Rosenblatt DS. Inborn errors of cobalamin absorption and metabolism. *Am J Med Genet C Semin Med Genet* 2011; 157C(1): 33–44. doi: 10.1002/ajmg.c.30288
47. Ozarda Y, Sikaris K, Streichert T, Macri J, IFCC Committee on Reference intervals and Decision Limits (C-RIDL). Distinguishing reference intervals and clinical decision limits – a review by the IFCC committee on reference intervals and decision limits. *Crit Rev Clin Lab Sci* 2018; 55(6): 420–31. doi: 10.1080/10408363.2018.1482256
48. Smith AD, Refsum H. Do we need to reconsider the desirable blood level of vitamin B12? *J Intern Med* 2012; 271(2): 179–82. doi: 10.1111/j.1365-2796.2011.02485.x
49. Vogiatzoglou A, Oulhaj A, Smith AD, Nurk E, Drevon CA, Ueland PM, et al. Determinants of plasma methylmalonic acid in a large population: implications for assessment of vitamin B12 status. *Clin Chem* 2009; 55(12): 2198–206. doi: 10.1373/clinchem.2009.128678
50. Fenech M. Folate (vitamin B9) and vitamin B12 and their function in the maintenance of nuclear and mitochondrial genome integrity. *Mutat Res* 2012; 733(1–2): 21–33. doi: 10.1016/j.mrfmmm.2011.11.003
51. Bjørke-Monsen AL. Assessment of cobalamin status. *Utredning av kobalaminstatus. Tidsskr Nor Laegeforen*. 2020; 140(9): 0589. doi: 10.4045/tidsskr.19.0589
52. Murphy MJ, Brandie F, Ebare M, Harrison M, Dow E, Bartlett WA, et al. Personalising laboratory medicine in the 'real world': assessing clinical utility, by clinical indication, of serum total B12 and active-B12(R) (holotranscobalamin) in the diagnosis of vitamin B12 deficiency. *Ann Clin Biochem* 2021; 58(5): 445–51. doi: 10.1177/00045632211003605
53. Hardlei TF, Morkbak AL, Bor MV, Bailey LB, Hvas AM, Nexø E. Assessment of vitamin B(12) absorption based on



- the accumulation of orally administered cyanocobalamin on transcobalamin. *Clin Chem* 2010; 56(3): 432–6. doi: 10.1373/clinchem.2009.131524
54. Bor MV, von Castel-Roberts KM, Kauwell GP, Stabler SP, Allen RH, Maneval DR, et al. Daily intake of 4 to 7 microg dietary vitamin B-12 is associated with steady concentrations of vitamin B-12-related biomarkers in a healthy young population. *Am J Clin Nutr* 2010; 91(3): 571–7. doi: 10.3945/ajcn.2009.28082
  55. Riedel BM, Molloy AM, Meyer K, Fredriksen A, Ulvik A, Schneede J, et al. Transcobalamin polymorphism 67A->G, but not 776C->G, affects serum holotranscobalamin in a cohort of healthy middle-aged men and women. *J Nutr* 2011; 141(10): 1784–90. doi: 10.3945/jn.111.141960
  56. Herrmann W, Obeid R, Schorr H, Geisel J. The usefulness of holotranscobalamin in predicting vitamin B12 status in different clinical settings. *Curr Drug Metab* 2005; 6(1): 47–53. doi: 10.2174/1389200052997384
  57. Murphy MM, Molloy AM, Ueland PM, Fernandez-Ballart JD, Schneede J, Arija V, et al. Longitudinal study of the effect of pregnancy on maternal and fetal cobalamin status in healthy women and their offspring. *J Nutr* 2007; 137(8): 1863–7. doi: 10.1093/jn/137.8.1863
  58. Molloy AM, Pangilinan F, Mills JL, Shane B, O'Neill MB, McGaughey DM, et al. A common polymorphism in HIBCH influences methylmalonic acid concentrations in blood independently of cobalamin. *Am J Hum Genet* 2016; 98(5): 869–82. doi: 10.1016/j.ajhg.2016.03.005
  59. Greibe E, Andreasen BH, Lildballe DL, Morkbak AL, Hvas AM, Nexø E. Uptake of cobalamin and markers of cobalamin status: a longitudinal study of healthy pregnant women. *Clin Chem Lab Med* 2011; 49(11): 1877–82. doi: 10.1515/CCLM.2011.682
  60. Lemming EW, Pitsi T. The Nordic Nutrition Recommendations 2022 - food consumption and nutrient intake in the adult population of the Nordic and Baltic countries. *Food Nutr Res*. 2022; 66: 8572. doi: 10.29219/fnr.v66.8572
  61. Davey GK, Spencer EA, Appleby PN, Allen NE, Knox KH, Key TJ. EPIC-Oxford: lifestyle characteristics and nutrient intakes in a cohort of 33 883 meat-eaters and 31 546 non meat-eaters in the UK. *Public Health Nutr* 2003; 6(3): 259–69. doi: 10.1079/PHN2002430
  62. Larsson CL, Johansson GK. Dietary intake and nutritional status of young vegans and omnivores in Sweden. *Am J Clin Nutr* 2002; 76: 100–6. doi: 10.1093/ajcn/76.1.100
  63. Leblanc JC, Yoon H, Kombadjian A, Verger P. Nutritional intakes of vegetarian populations in France. *Eur J Clin Nutr* 2000; 54(5): 443–9. doi: 10.1038/sj.ejcn.1600967
  64. Neufingerl N, Eilander A. Nutrient Intake and Status in Adults Consuming Plant-Based Diets Compared to Meat-Eaters: A Systematic Review. *Nutrients*. 2021; 14(1): 29. doi: 10.3390/nu14010029
  65. Zugravu CA, Macri A, Belc N, Bohiltea R. Efficacy of supplementation with methylcobalamin and cyanocobalamin in maintaining the level of serum holotranscobalamin in a group of plant-based diet (vegan) adults. *Exp Ther Med* 2021; 22(3): 993. doi: 10.3892/etm.2021.10425
  66. Bryngelsson S, Moshtaghian H, Bianchi M, Hallstrom E. Nutritional assessment of plant-based meat analogues on the Swedish market. *Int J Food Sci Nutr* 2022; 73(7): 889–901. doi: 10.1080/09637486.2022.2078286
  67. Henjum S, Grouffh-Jacobsen S, Allen LH, Raael E, Israelsson AM, Shahab-Ferdows S, et al. Adequate vitamin B12 and folate status of Norwegian vegans and vegetarians [published online ahead of print, 2022 Sep 27]. *Br J Nutr*. 2022; 129(12): 1–8. doi: 10.1017/S0007114522002987
  68. Greibe E, Nexø E. Forms and amounts of vitamin B12 in infant formula: a pilot study. *PLoS One* 2016; 11(11): e0165458. doi: 10.1371/journal.pone.0165458
  69. Ford C, Rendle M, Tracy M, Richardson V, Ford H. Vitamin B12 levels in human milk during the first nine months of lactation. *Int J Vitam Nutr Res* 1996; 66(4): 329–31.
  70. Henjum S, Manger M, Hampel D, Brantsæter AL, Shahab-Ferdows S, Bastani NE, et al. Vitamin B12 concentrations in milk from Norwegian women during the six first months of lactation. *Eur J Clin Nutr*. 2020; 74(5): 749–756. doi: 10.1038/s41430-020-0567-x. Epub 2020 Jan 30. PMID: 32001810; PMCID: PMC7214247.
  71. Elizabeth KE, Krishnan V, Vijayakumar T. Umbilical cord blood nutrients in low birth weight babies in relation to birth weight & gestational age. *Indian J Med Res* 2008; 128(2): 128–33.
  72. Lindenbaum J, Healton EB, Savage DG, Brust JC, Garrett TJ, Podell ER, et al. Neuropsychiatric disorders caused by cobalamin deficiency in the absence of anemia or macrocytosis. *N Engl J Med* 1988; 318(26): 1720–8. doi: 10.1056/NEJM198806303182604
  73. Dharmarajan TS, Adiga GU, Norkus EP. Vitamin B12 deficiency. Recognizing subtle symptoms in older adults. *Geriatrics* 2003; 58(3): 30–4, 7–8.
  74. Smith AD, Warren MJ, Refsum H. Vitamin B12. *Adv Food Nutr Res* 2018; 83: 215–79. doi: 10.1016/bs.afnr.2017.11.005
  75. Bennett M. Vitamin B12 deficiency, infertility and recurrent fetal loss. *J Reprod Med* 2001; 46(3): 209–12.
  76. Bondevik GT, Schneede J, Refsum H, Lie RT, Ulstein M, Kvale G. Homocysteine and methylmalonic acid levels in pregnant Nepali women. Should cobalamin supplementation be considered? *Eur J Clin Nutr* 2001; 55(10): 856–64. doi: 10.1038/sj.ejcn.1601236
  77. Kirke PN, Molloy AM, Daly LE, Burke H, Weir DG, Scott JM. Maternal plasma folate and vitamin B12 are independent risk factors for neural tube defects. *Q J Med* 1993; 86(11): 703–8.
  78. Banihani SA. Vitamin B12 and Semen Quality. *Biomolecules*. 2017; 7(2): 42. doi: 10.3390/biom7020042
  79. Holland D, Chang L, Ernst TM, Curran M, Buchthal SD, Alicata D, et al. Structural growth trajectories and rates of change in the first 3 months of infant brain development. *JAMA Neurol* 2014; 71(10): 1266–74. doi: 10.1001/jamaneurol.2014.1638
  80. Larnkjer A, Christensen SH, Lind MV, Michaelsen KF, Mølgaard C. Plasma vitamin B12 concentration is positively associated with cognitive development in healthy Danish 3-year-old children: the SKOT cohort studies. *Br J Nutr*. 2022; 128(10): 1946–1954. doi: 10.1017/S0007114521004888
  81. Kang D, Shi B, Erfe MC, Craft N, Li H. Vitamin B12 modulates the transcriptome of the skin microbiota in acne pathogenesis. *Sci Transl Med* 2015; 7(293): 293ra103. doi: 10.1126/scitranslmed.aab2009
  82. Bjørke-Monsen AL, Ulvik A, Nilsen RM, Midttun Ø, Roth C, Magnus P, et al. Impact of Pre-Pregnancy BMI on B Vitamin and Inflammatory Status in Early Pregnancy: An Observational Cohort Study. *Nutrients* 2016; 8(12): 776. doi: 10.3390/nu8120776. PMID: 27916904; PMCID: PMC5188431
  83. Collin SM, Metcalfe C, Refsum H, Lewis SJ, Zuccolo L, Smith GD, et al. Circulating folate, vitamin B12, homocysteine, vitamin B12 transport proteins, and risk of prostate cancer: a case-control study, systematic review, and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2010; 19(6): 1632–42. doi: 10.1158/1055-9965.EPI-10-0180

84. Arendt JFH, Sorensen HT, Horsfall LJ, Petersen I. Elevated vitamin B12 levels and cancer risk in UK primary care: a THIN database cohort study. *Cancer Epidemiol Biomarkers Prev* 2019; 28(4): 814–21. doi: 10.1158/1055-9965.EPI-17-1136
85. Arendt JF, Pedersen L, Nexo E, Sorensen HT. Elevated plasma vitamin B12 levels as a marker for cancer: a population-based cohort study. *J Natl Cancer Inst* 2013; 105(23): 1799–805. doi: 10.1093/jnci/djt315
86. Arendt JF, Nexo E. Cobalamin related parameters and disease patterns in patients with increased serum cobalamin levels. *PLoS One* 2012; 7(9): e45979. doi: 10.1371/journal.pone.0045979
87. Andreeva VA, Touvier M, Kesse-Guyot E, Julia C, Galan P, Hercberg S. B vitamin and/or omega-3 fatty acid supplementation and cancer: ancillary findings from the supplementation with folate, vitamins B6 and B12, and/or omega-3 fatty acids (SU.FOL.OM3) randomized trial. *Arch Intern Med* 2012; 172(7): 540–7. doi: 10.1001/archinternmed.2011.1450
88. Zhang SL, Chen TS, Ma CY, Meng YB, Zhang YF, Chen YW, et al. Effect of vitamin B supplementation on cancer incidence, death due to cancer, and total mortality: a PRISMA-compliant cumulative meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2016; 95(31): e3485. doi: 10.1097/MD.0000000000003485
89. Zhang SM, Cook NR, Albert CM, Gaziano JM, Buring JE, Manson JE. Effect of combined folic acid, vitamin B6, and vitamin B12 on cancer risk in women: a randomized trial. *JAMA* 2008; 300(17): 2012–21. doi: 10.1001/jama.2008.555
90. Sun NH, Huang XZ, Wang SB, Li Y, Wang LY, Wang HC, et al. A dose-response meta-analysis reveals an association between vitamin B12 and colorectal cancer risk. *Public Health Nutr* 2016; 19(8): 1446–56. doi: 10.1017/S136898001500261X
91. Moore E, Mander A, Ames D, Carne R, Sanders K, Watters D. Cognitive impairment and vitamin B12: a review. *Int Psychogeriatr* 2012; 24(4): 541–56. doi: 10.1017/S1041610211002511
92. Malouf R, Areosa Sastre A. Vitamin B12 for cognition. *Cochrane Database Syst Rev* 2003; (3): CD004326. doi: 10.1002/14651858.CD004326
93. Smith AD. The worldwide challenge of the dementias: a role for B vitamins and homocysteine? *Food Nutr Bull* 2008; 29(2 Suppl): S143–72. doi: 10.1177/15648265080292S119
94. Douauid G, Refsum H, de Jager CA, Jacoby R, Nichols TE, Smith SM, et al. Preventing Alzheimer's disease-related gray matter atrophy by B-vitamin treatment. *Proc Natl Acad Sci U S A* 2013; 110(23): 9523–8. doi: 10.1073/pnas.1301816110
95. Ma F, Zhou X, Li Q, Zhao J, Song A, An P, et al. Effects of folic acid and vitamin B12, alone and in combination on cognitive function and inflammatory factors in the elderly with mild cognitive impairment: a single-blind experimental design. *Curr Alzheimer Res* 2019; 16(7): 622–32. doi: 10.2174/1567205016666190725144629
96. Kim JM, Stewart R, Kim SW, Yang SJ, Shin IS, Yoon JS. Predictive value of folate, vitamin B12 and homocysteine levels in late-life depression. *Br J Psychiatry* 2008; 192(4): 268–74. doi: 10.1192/bjp.bp.107.039511
97. Sangle P, Sandhu O, Aftab Z, Anthony AT, Khan S. Vitamin B12 supplementation: preventing onset and improving prognosis of depression. *Cureus* 2020; 12(10): e11169. doi: 10.7759/cureus.11169
98. Almeida OP, Ford AH, Flicker L. Systematic review and meta-analysis of randomized placebo-controlled trials of folate and vitamin B12 for depression. *Int Psychogeriatr* 2015; 27(5): 727–37. doi: 10.1017/S1041610215000046
99. Darby WJ, Bridgforth EB, Le Brocqy J, Clark SL, Jr., De Oliveira JD, Kevany J, et al. Vitamin B12 requirement of adult man. *Am J Med* 1958; 25(5): 726–32. doi: 10.1016/0002-9343(58)90010-X
100. Herbert V. Vitamin B12: plant sources, requirements, and assay. *Am J Clin Nutr* 1988; 48: 852–8. doi: 10.1093/ajcn/48.3.852
101. EFSA. EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA). Scientific Opinion on Dietary Reference Values for cobalamin (vitamin B12) 2015. *EFSA Journal* 13: 4150. Available from: <file:///C:/Temp/j.efsa.2015.4150.pdf>. accessed 04 March 2022.
102. Bor MV, Lydeking-Olsen E, Moller J, Nexo E. A daily intake of approximately 6 microg vitamin B-12 appears to saturate all the vitamin B-12-related variables in Danish postmenopausal women. *Am J Clin Nutr* 2006; 83(1): 52–8. doi: 10.1093/ajcn/83.1.52
103. Vogiatzoglou A, Smith AD, Nurk E, Berstad P, Drevon CA, Ueland PM, et al. Dietary sources of vitamin B-12 and their association with plasma vitamin B-12 concentrations in the general population: the Hordaland Homocysteine Study. *Am J Clin Nutr* 2009; 89(4): 1078–87. doi: 10.3945/ajcn.2008.26598
104. Pellinen T, Paivarinta E, Isotalo J, Lehtovirta M, Itonen ST, Korkalo L, et al. Replacing dietary animal-source proteins with plant-source proteins changes dietary intake and status of vitamins and minerals in healthy adults: a 12-week randomized controlled trial. *Eur J Nutr* 2022; 61(3): 1391–404. doi: 10.1007/s00394-021-02729-3
105. Schorah CJ, Smithells RW, Scott J. Vitamin B12 and anencephaly. *Lancet* 1980; 1: 880. doi: 10.1016/S0140-6736(80)91381-1
106. Bjørke-Monsen AL, Roth C, Magnus P, Midttun O, Nilsen RM, Reichborn-Kjennerud T, et al. Maternal B vitamin status in pregnancy week 18 according to reported use of folic acid supplements. *Mol Nutr Food Res* 2013; 57(4): 645–52. doi: 10.1002/mnfr.201200114
107. Fokkema MR, Wolt HA, van Beusekom CM, Schaafsma A, Dijk-Brouwer DA, Muskiet FA. Plasma total homocysteine increases from day 20 to 40 in breastfed but not formula-fed low-birthweight infants. *Acta Paediatr* 2002; 91(5): 507–11. doi: 10.1111/j.1651-2227.2002.tb03268.x
108. Hay G, Trygg K, Whitelaw A, Johnston C, Refsum H. Folate and cobalamin status in relation to diet in healthy 2-y-old children. *Am J Clin Nutr* 2011; 93(4): 727–35. doi: 10.3945/ajcn.110.003426
109. Folkehelseinstituttet. UNGKOST 3 Landsomfattende kostholdsundersøkelse blant elever i 4. -og 8. klasse i. 2015. Available from: <https://www.fhi.no/globalassets/dokumenterfiler/rapporter/2016/ungkost-rapport-24.06.16.pdf>. accessed 04 March 2022.
110. Bjørke Monsen AL, Ueland PM. Homocysteine and methylmalonic acid in diagnosis and risk assessment from infancy to adolescence. *Am J Clin Nutr* 2003; 78(1): 7–21. doi: 10.1093/ajcn/78.1.7
111. MacFarlane AJ, Shi Y, Greene-Finestone LS. High-dose compared with low-dose vitamin B-12 supplement use is not associated with higher vitamin B-12 status in children, adolescents, and older adults. *J Nutr* 2014; 144(6): 915–20. doi: 10.3945/jn.113.190256

---

**\*Anne-Lise Bjørke Monsen**

Department of Medical Biochemistry and Pharmacology  
Haukeland University Hospital  
Bergen, Norway  
Email: [almo@helse-bergen.no](mailto:almo@helse-bergen.no)